

The Art of Error Correcting Coding  
Robert H. Morelos-Zaragoza  
Second Edition, John Wiley & Sons, 2006  
ISBN: 0470015586

Errata (Updated 12/4/2007)

Chapter 1

Page 13: In equation (1.27) the limit of the summation should be  $t$  and not  $n$ :

$$\sum_{i=0}^t \binom{n}{i} (q-1)^i \leq q^{n-k}.$$

Chapter 3

Page 70: The parameters of the BCH codes in problems 8 and 9 need to be exchanged. Namely, (15, 7, 5) for problem 8 and (15, 5, 7) for problem 9, as follows:

8. Show that the (0,2)-order EG code of length 15 is a binary BCH (15, 7, 5) code.
9. Show that the (1,1)-order EG code of length 15 is a binary BCH (15, 5, 7) code.

In problem 14, part (a), the problem is number 6 (not 2).

Chapter 4

Page 78: In Equation (4.7), the argument of  $z(x)$  the exponent is negative. (The inverse of the position, eq. (10.14) of Berlekamp (1984)):

$$e_{j\ell} = \frac{(\alpha^{j\ell})^{1-b} z(\alpha^{-j\ell})}{\nu \prod_{\substack{i=1 \\ i \neq \ell}} (\alpha^{j\ell})^{i-j\ell}}$$

Page 85: In problem 4 and 5, the primitive polynomials are  $p(x) = x^3 + x + 1$  and  $p(x) = x^5 + x^2 + 1$ , respectively.

Chapter 5

Page 117: In problem 8, the free distance is 3. For problem 9, use the most significant term of the polynomial in (5.20). This values of this term are  $\{3x^6, 42x^5, 92x^4\}$  for rates  $\{2/3, 3/4, 5/6\}$ , respectively. (Appendices A and C of (Lee 1997)).


Chapter 6

Page 141: In problem 9 use a “ $|u|v + w|$ ” construction (and not construction X), with  $u, v$  and  $w$  denoting codewords in  $C_1, C_2$  and  $C_3$  respectively.

Chapter 7

Page 160: Figure 7.11 is not correct and needs to be modified as do the soft outputs (one per information bit, excluding the tail bits) which should be: +18,+14,-4,-2. The correct figure is shown below.

Transmitted	-1,-1	+1,-1	+1,-1	+1,+1	-1,+1	-1,-1
Received	-4,-1	-1,-3	+2,-3	+3,+3	-3,+3	+3,+1



Page 162: The subindex of  $\gamma$  in (7.25) is  $i + 1$  and its argument  $(m, m')$ , as in (7) of Bahl et al. (1974):

$$\beta_i(m) = \sum_{m'} \beta_{i+1}(m') \cdot \sum_{j=0}^1 \gamma_{i+1}^{(j)}(m, m')$$

Page 168: Assume that  $N_0 = 10$  in problem 9. In problem 10, by zero-tail it is meant that all trellis paths terminate at the all-zero state.

Chapter 8

Page 201: In problem 2, you may approximate the LLR value of a parity-check bit as "the negative of the product of the signs times the minimum amplitude (reliability)" similar to (12) of (Hagenauer et al. 1996).

$$\Lambda(x_1 \oplus x_2 \oplus \dots \oplus x_\ell) \approx - \left[ \prod_{j=1}^{\ell} (-1)^{\text{sgn}(\Lambda(x_j))} \right] \cdot \min_{1 \leq j \leq \ell} \{ |\Lambda(x_j)| \}.$$

Page 202: The function in question is  $G(x) = \log(F(x))$  and not  $F(x)$ . Prove that  $G(G(x)) = x$ .

Chapter 9

Page 212: The values in Figure 9.12 should be the squared Euclidean distances:  $a = 4$ ,  $b = 0.586$ ,  $c = 2$  and  $d = 3.414$ .